

# Forward $\pi^0$ and $\eta$ production in STAR at $\sqrt{s}=500$ GeV with transversely polarized pp collisions

## *Transverse momentum Dependence of $\pi^0$ SSA in FMS Run 11 CIPANP*

S. Heppelmann (PSU) for STAR collaboration  
June 2, 2012

- Background
  - Physics Questions
  - **Cross Ratio** method vs.  $A(\phi)=A_N \cos(\phi)$  fitting method
  - Previous FMS and STAR results
  - About  $P_T$  dependence of  $A_N$
  - FMS Event Topology and Event Selection
- Present High Statistics  $A_N$  for STAR Run 11  $\sqrt{s}=500$  GeV
  - $X_F$  dependence
  - $P_T$  dependence for fixed  $X_F$
  - Dependence on event topology



# Proton Forward Scattering at High PT


## QCD Perspective

### PQCD (Leading Twist):

Factorized Cross Section= (initial state) x (quark scattering) x (fragmentation)


- Does good job of predicting the spin averaged cross section.

- Leading twist cross section does not depend on transverse polarization.
- Spin Dependence require refinements like:
  - Beyond Collinear Factorization (**Sivers**)
  - Models of spin dependent factorization (**Collins**)
  - Models that go **beyond leading twist**.



## FMS

Pb Glass EM Calorimeter  
pseudo-rapidity  $2.7 < \eta < 4.0$   
Small cells:  $3.81 \times 3.81$  cm  
Outer cells:  $5.81 \times 5.81$  cm




Unpolarized  
Proton


$\pi^0$



Transversely  
Polarized  
Proton



**FPD EM Calorimeter**  
Small cells only  
Two  $7 \times 7$  arrays



**Forward EM Calorimetry In STAR.**


# 1) Cross Ratio Transverse Asymmetry

VS


# 2) $A(\phi)$ Fit

**Method 1:**  
**Cross Ratio:**

$$A_N = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow} \cong \frac{1}{P} \frac{\sqrt{S^\uparrow N^\downarrow} - \sqrt{N^\uparrow S^\downarrow}}{\sqrt{N^\uparrow S^\downarrow} + \sqrt{S^\uparrow N^\downarrow}}$$



Left(S):  $\cos(\phi) > 0.5$



Viewed from  
collision point

Right(N):  $\cos(\phi) < -0.5$

**Method 2:**  $a_N(\phi) = a_0 + A_N \cos(\phi)$

**Fix  $a_0$  for full data set**



**For many small data subsets ..... one parameter fit for  $A_N$**

**Advantage: Every fitted value of  $A_N$  comes with error and  $\chi^2$ .**




# New paper on $\eta/\pi^0$ at $X_F > 0.5$

Preprint Ref to be added here!





- $\pi^0$  cross section in **good agreement with PQCD calculation.**
- $\eta/\pi^0$  cross section ratio similar to that observed where jet fragmentation is dominant.
- $A_N(\eta) > A_N(\pi^0)$  for  $X_F > 0.55$



# STAR Published Run 6 (FPD $\sqrt{s}=200\text{GeV}$ )

PRL 101, 222001 (2006)


- Rising  $A_N$  with  $X_F$  ( $0 < X_F < 0.5$ ) from 0% to 5-10%
- No evidence of fall in  $A_N$  with increasing  $P_T$  up to  $P_T \sim 3 \text{ GeV}/c$





# From FMS **Run 8**, STAR has Expanded Rapidity Coverage $-1 < Y < 4.2$

**STAR** Forward Meson Spectrometer  
 $2.5 < Y < 4.0$



**arXiv:0901.2763 +**  
**A.Ogawa @CIPANP09**



- Leading twist cross section does not depend on transverse polarization.
- Spin Dependence require refinements like:
  - Beyond Collinear Factorization (**Sivers**)
  - Models of spin dependent factorization (**Collins**)
  - Models that go **beyond leading twist**.

**Sivers Model:** Initial quark picks up  $k_T$  from initial state wave function, **proportional to orbital angular momentum**.

Jet based Asymmetry, significant dependence of  $A_N$  on the details of near side jet fragments is not expected!

**Collins Model:** Final  $\pi^0$  picks up  $k_T$  from **fragmentation of polarized quark**. Vanishing jet asymmetry. Observed  $A_N$  will depend on the details of near side fragmentation!

A toy model for proton  
Cross Section at large  $x$ .

$$\sigma(p_T) \sim \frac{(1-x_F)^5}{p_T^6}$$

**Suppose** initial state structure or final state fragmentation modifies the hard scattering  $\mathbf{p}_T$ .

If the spin dependent initial/final state momentum is  $\mathbf{k}_T$ .

For spin proton spin up:  $\langle \mathbf{p}_T \rangle \Rightarrow \langle \mathbf{p}_T \rangle - \mathbf{k}_T$

For spin proton spin dn:  $\langle \mathbf{p}_T \rangle \Rightarrow \langle \mathbf{p}_T \rangle + \mathbf{k}_T$

$$A_N(p_T) \sim \frac{\sigma(p_T - k_T) - \sigma(p_T + k_T)}{2\sigma(p_T)} \sim \frac{-k_T}{\sigma} \frac{d\sigma}{dp_T} \sim \frac{6k_T}{p_T} \propto \frac{1}{p_T}$$

Similar for **for higher twist:**



$$A_N(p_T) \propto \frac{1}{p_T}$$



# Isolation of $\pi^0$ 's



## Event Selection:

1. **Analyze FMS for all photon** candidates.  
(Showers that are fit successfully to photon hypothesis)  
A photon candidates must have a minimum of 6 GeV in the small inner detector or 4 GeV in the outer cells.
2. **Find Clusters of EM energy** grouping photon candidates that are within opening angle cone  $\Delta\theta$  (relative to energy weighted center)
3. We consider 2 event classes {1 and 2}
  1.  $\Delta\theta = 0.07$  2 Photon clusters,  $\pi^0$  Mass (isolation radius of .07 radians).
  2.  $\Delta\theta = 0.03$  2 Photon clusters,  $\pi^0$  Mass (isolation radius of .03 radians).



## Class 1 Events: $\Delta\theta = 0.07$ 2 Photon clusters, $\pi^0$ Mass (less inclusive)?

- $40 \text{ GeV} < E_{\text{pair}} < 100 \text{ GeV}$
- $Z = |(E_1 - E_2)/(E_1 + E_2)| < 0.7$
- $2.7 < Y < 4.0$  (Full FMS Pseudo-rapidity)
- Selection of  $\pi^0$  Peak ( $0.02 < \text{Mass} < 0.3$ )
- Average polarization:  $51.6\% \pm 6.7\%$  (RHIC Spin CNI Group <http://www.phy.bnl.gov/cnipol/>)
- Integrated Luminosity:  $22 \text{ pb}^{-1}$





STAR Preliminary

# Cross Ratio Transverse Single Spin Asymmetry for Run 11


$\pi^0$  (2 Photon Cluster) **Cluster size = 0.07 Rad**

For **Blue** Beam (Forward)

Full FMS rapidity range ( $2.6 < Y < 4.1$ )





Left(S):  $\cos(\phi) > 0.5$



Right(N):  $\cos(\phi) < -0.5$





Compare **new  $\sqrt{s}=500$  GeV Run 11 Full FMS Data** on right with **Run 6  $\sqrt{s}=200$**  published data below.




Compare **new  $\sqrt{s}=500$  GeV Run 11** Full FMS Data on right with **Run 6  $\sqrt{s}=200$**  published data below.

Scale of  $A_N$  similar but starts at lower  $X_F$  in Run 11 data.







Blue Beam  $A_N$


As an alternative to Cross Ratio, the raw asymmetry can be plotted as a function of  $\cos(\phi)$  (with polarization axis at  $\phi = \pi/2$ )

Slope =  $A_N$



Intercept = Luminosity Ratio for data set

Luminosity ratio for all  $\sim -0.31 \pm 0.05 \%$



Slope Fits are consistent with Cross Ratio Method.





# Transverse Single Spin $\pi^0$ Asymmetry vs $P_T$ for small and large $\pi^0$ isolation cones. (Errors shown are statistical)



# Transverse Single Spin $\pi^0$ Asymmetry vs $P_T$ for small and large $\pi^0$ isolation cones. (Errors shown are statistical)



# Transverse Single Spin $\pi^0$ Asymmetry vs $P_T$ for small and large $\pi^0$ isolation cones. (Errors shown are statistical)






# **Transverse Single Spin $\pi^0$ Asymmetry vs $P_T$ for small and large $\pi^0$ isolation cones.** (Errors shown are statistical)

Higher Twist or other pQCD related models suggest  $A_N$  should fall at large  $P_T$  with at least 1 power of  $P_T$

These plots include 2 parameter fits for  $A_N$  vs  $P_T$ :

$$A_N(P_T) = [p_0] \times (P_T)^{[p_1]}$$

Fits are shown for both the **70 mRad** and **30 mRad** isolation cones.





# Systematic Errors

- Run 11 blue beam polarization  $51.6\% \pm 6.7$
- Non  $\pi^0$  signal  $< 10\%$
- Similar asymmetries for Background:

$$\frac{\Delta A_N}{A_N} < 13\%$$

$$\frac{\Delta A_N}{A_N} < 5\%$$

$$\frac{\Delta P_T}{P_T} < 12\%$$

$$\frac{\Delta A_N}{A_N} < 5\%$$

- $P_T$  uncertainty
  - Energy 10%
  - Angle 6%

$$\frac{\Delta P_T}{P_T} < 12\%$$

$$\frac{\Delta A_N}{A_N} < 5\%$$

Total Systematic Asymmetry Error  
Common to all data points.

$$\frac{\Delta A_N}{A_N} < 15\%$$



# Conclusion

## STAR $\pi^0$ $A_N$ at $\sqrt{s}=500$ GeV

- $A_N$  increases with  $X_F$  (as seen at lower energies).
- $A_N$  less dependent on  $P_T$  than models predict to  $P_T \sim 10$  GeV/c. Data may be consistent with flat dependence on  $P_T$ .
- For data points at  $X_F < 0.32$ ,  $A_N$  is significantly larger when the  $\pi^0$ s are more isolated (0.07 Rad).

Additional E&M signals in the same general direction as the  $\pi^0$  ( $> \sim 5$  GeV between 0.03 and 0.07 radians from the  $\pi^0$ )  
contribute little to the observed Transverse Single Spin Asymmetry.

- **New Data Coming RHIC RUN 12**
  - ~20 pb<sup>-1</sup> of  $\sqrt{s}=200$  GeV pp
  - ~Transversely Polarized FMS data
  - ~ Similar measurement up to  $P_T > 6$  GeV/c

