

**Measurement of W^\pm single spin asymmetries in polarized $p + p$ collisions at
 $\sqrt{s} = 510$ GeV at STAR**

Devika Gunarathne for the STAR Collaboration

Temple University, Philadelphia, PA, USA

Email: devika.gunarathne@temple.edu

The STAR experiment at RHIC has provided significant contributions to our understanding of the structure of the proton. The STAR experiment is well equipped to measure $W^\pm \rightarrow e^\pm + \nu$ in $\sqrt{s} = 510$ GeV longitudinally polarized $p + p$ collisions. The longitudinal single spin asymmetry in W production, A_L , measured as a function of decay positron (electron) pseudo-rapidity η for W^+ (W^-) is sensitive to the individual helicity polarizations of u and \bar{d} (d and \bar{u}) quarks. Due to maximal violation of parity during the production, W bosons couple to left-handed quarks and right-handed anti-quarks and hence offer direct probes of their respective helicity distributions in the nucleon. The published STAR A_L results (combination of 2011 and 2012 data) have been used by several theoretical analyses suggesting a significant impact in constraining the helicity distributions of \bar{u} , and \bar{d} quarks. In 2013 STAR collected a dataset at $\sqrt{s} = 510$ GeV with a total integrated luminosity of ~ 300 pb $^{-1}$ with an average beam polarization of $\sim 54\%$, a figure of merit three times larger than the dataset used by previous analyses. We will present preliminary results of STAR 2013 W A_L measurement at mid-rapidity ($|\eta| < 1$) region.